

A simple realization of $gl(2, c)$ Lie algebra with vector operators on sphere

Q. H. Liu,¹ X. P. Rong,¹ and D. M. Xun¹

¹*School for Theoretical Physics, and Department of Applied Physics, Hunan University, Changsha, 410082, China*

(Dated: January 29, 2010)

By utilization of three elementary vector operators as position, angular momentum and their cross product, a simple realization of $gl(2, c)$ Lie algebra on sphere are constructed. The coherent states based on this algebra can then be constructed by standard manner.

PACS numbers: 03.65.Ca

I. INTRODUCTION

Position operator $\mathbf{R}$, angular momentum operator $\mathbf{L}$ and their cross product as $\mathbf{R} \times \mathbf{L}$ are three typical vector operators [1, 2] that are widely used to explore the quantum properties of the angular momentum [3], and various coherent states on the spheres [4–7]. However, in contrast with the former two operators that are well-studied, the fundamental properties of the operator $\mathbf{R} \times \mathbf{L}$ are still not fully understood yet. In this paper, we present an algebraic analysis of the linear combination of three vector operators of following form,

$$\mathbf{J} = \frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + a \mathbf{N} + \frac{b}{\hbar} \mathbf{N} L_z, \quad (1)$$

where $\mathbf{N} \equiv \mathbf{R}/R$ is the unit directional vector, and a and b are two parameters. We are going to point out that, with proper combinations, this vector (1) together with $\mathbf{L}$ can be used to form the $gl(2, c)$ Lie algebra. As well-known, the $gl(2, c)$ algebra is in general the four dimensional Lie algebra $G(a_0, b_0)$ with a_0 and b_0 being two parameters, composed of operators K_+ , K_- , and K_z and I ; and these operators satisfy commutation relations [8]:

$$\begin{cases} [K_+, K_-] = 2a_0 K_z + b_0 I \\ [K_z, K_{\pm}] = \pm K_{\pm} \\ [\mathbf{K}, I] = 0 \end{cases}, \quad (2)$$

where I is a c -number, $\mathbf{K} = (K_x, K_y, K_z)$ and $K_{\pm} = K_x \pm iK_y$ as usual. Two Casimir operators are [8]:

$$\begin{cases} C = K_+ K_- + a_0 K_z^2 - (a_0 + b_0) K_z \\ C_0 = I \end{cases}. \quad (3)$$

As a_0 and b_0 take values 0 or ± 1 , one has the following algebras [8–11]:

$$G(1, \pm 1) : o(3) \oplus u(1) \approx u(2) \oplus u(1), \quad (4)$$

$$G(-1, \pm 1) : o(2, 1) \oplus u(1) \approx u(1, 1) \oplus u(1), \quad (5)$$

$$G(1, 0) : so(3) \oplus u(1) \approx su(2) \oplus u(1), \quad (6)$$

$$G(-1, 0) : so(2, 1) \oplus u(1) \approx su(1, 1) \oplus u(1), \quad (7)$$

and etc [8–11].

This paper is organized as what follows. In section II, the raising operator K_+ and the lowering operator K_- are formed by the proper construction of vectors $\mathbf{J}$ (1). In section III, an algebraic analysis is performed. In last section IV, a brief conclusion is given.

II. RAISING OPERATOR K_+ AND LOWING OPERATOR K_-

For convenience, we write down two operators of similar form,

$$\begin{cases} \mathbf{J}_1 = \frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + a \mathbf{N} + \frac{b}{\hbar} \mathbf{N} L_z \\ \mathbf{J}_2 = -\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + c \mathbf{N} + \frac{d}{\hbar} \mathbf{N} L_z \end{cases}, \quad (8)$$

where c and d are new parameters similar to a and b . The hermitian conjugate of these two operators $\mathbf{J}_1$ and $\mathbf{J}_2$ are given by respectively,

$$\begin{cases} \mathbf{J}_1^\dagger = -\frac{i}{\hbar}\mathbf{N} \times \mathbf{L} + a^*\mathbf{N} + \frac{b^*}{\hbar}\mathbf{N}L_z - 2\mathbf{N} + \frac{b^*}{\hbar}[L_z, \mathbf{N}] \\ \mathbf{J}_2^\dagger = \frac{i}{\hbar}\mathbf{N} \times \mathbf{L} + c^*\mathbf{N} + \frac{d^*}{\hbar}\mathbf{N}L_z + 2\mathbf{N} + \frac{d^*}{\hbar}[L_z, \mathbf{N}] \end{cases} \quad (9)$$

Now, we define raising operator K_+ and lowering operator K_- by $K_+ \equiv J_{1x} + iJ_{1y}$ and $K_- \equiv J_{2x} - iJ_{2y}$,

$$K_+ = \frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_+ + aN_+ + bN_+ \frac{L_z}{\hbar}, \quad (10)$$

$$K_- = -\frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_- + cN_- + dN_- \frac{L_z}{\hbar}. \quad (11)$$

Their hermitian conjugates are after calculations,

$$K_+^\dagger = -\frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_- + (a^* - 2 - b^*)N_- + b^*N_- \frac{L_z}{\hbar}, \quad (12)$$

$$K_-^\dagger = \frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_+ + (c^* + 2 + d^*)N_+ + d^*N_+ \frac{L_z}{\hbar}. \quad (13)$$

We impose following requirement $K_+^\dagger = K_-$ on these operators, and find,

$$\begin{cases} a^* - 2 - b^* = c \\ b^* = d \end{cases} \quad (14)$$

Taking the hermitian conjugate twice, we will get back to the same operator $K_-^\dagger = (K_+^\dagger)^\dagger = K_+$; and the results identical to Eq. (14) will be obtained. For further simplification, we require that these parameters a , b , c and d are all real numbers. In consequence, there are only two parameters are independent, which can be conveniently chosen as a and b ,

$$c = a - b - 2, d = b. \quad (15)$$

Finally we get the the raising and the lowering operator K_+ and K_- respectively,

$$K_+ = \frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_+ + aN_+ + bN_+ \frac{L_z}{\hbar}, \text{ and } K_- = -\frac{i}{\hbar}(\mathbf{N} \times \mathbf{L})_- + (a - b - 2)N_- + bN_- \frac{L_z}{\hbar}. \quad (16)$$

The third operator K_z is not known yet, which will be determined by requiring that that three operators (K_x, K_y, K_z) form the $gl(2, c)$ Lie algebra through the communication relations (2). The explicit form of K_z will be carried out in following section.

III. LIE ALGEBRA ANALYSIS OF K_+ AND K_-

Direct computation can give following relations,

$$[K_+, K_-] = -(a + c)(b + 1) - ((a + c)(1 - b))N_z^2 - 2(1 + 2b + b^2 + b(1 - b)N_z^2) \frac{L_z}{\hbar}, \quad (17)$$

$$[L_z/\hbar, K_\pm] = \pm K_\pm. \quad (18)$$

These two relations suffice to meet our need, though there are still other communication relations. The closure of the $gl(2, c)$ Lie algebra implies that two coefficients before N_z^2 are zero,

$$\begin{cases} (a + c)(1 - b) = 0 \\ b(1 - b) = 0 \end{cases} \quad (19)$$

Thus, with redefining operator $K_z \equiv L_z/\hbar$, we finally obtain the $gl(2, c)$ Lie algebra:

$$[K_+, K_-] = -(a + c)(b + 1) - 2(b + 1)^2 K_z, \quad [K_z, K_\pm] = \pm K_\pm. \quad (20)$$

All possibilities of the independent parameters a and b in Eq. (20) are given by association of Eqs. (15) and (19)

$$\begin{cases} i) & b = 0, a = 1 \\ ii) & b = 1, a \neq 1 \\ iii) & b = 1, a = 1 \end{cases} . \quad (21)$$

So, we have only three decompositions of the $gl(2, c)$ Lie algebra: Case 1, $su(1, 1)$ with $a = -c = 1$ and $b = d = 0$, case 2, $u(1, 1) \otimes u(1)$ with $b = d = 1$ and $a + c \neq 0$, and finally case 3, $su(1, 1)$ with $b = d = 1$ and $a = -c = 1$. In all these cases, $K_z \equiv L_z/\hbar$, and for case 1, 2, and 3, communication relations $[K_+, K_-]$ are given by, respectively,

$$\begin{cases} K_x = i \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \mathbf{N} \right)_y \\ K_y = -i \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \mathbf{N} \right)_x \end{cases} , \quad (22)$$

$$\begin{cases} K_x = \left(\frac{(a+c)}{2} \mathbf{N} + \frac{1}{\hbar} \mathbf{N} L_z \right)_x + i \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \frac{(a-c)}{2} \mathbf{N} \right)_y \\ K_y = -i \left\{ \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \frac{(a-c)}{2} \mathbf{N} \right)_x + i \left(\frac{(a+c)}{2} \mathbf{N} + \frac{1}{\hbar} \mathbf{N} L_z \right)_y \right\} \end{cases} , \quad (23)$$

$$\begin{cases} K_x = \left(\frac{1}{\hbar} \mathbf{N} L_z \right)_x + i \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \mathbf{N} \right)_y \\ K_y = -i \left(\frac{i}{\hbar} \mathbf{N} \times \mathbf{L} + \mathbf{N} \right)_x + \left(\frac{1}{\hbar} \mathbf{N} L_z \right)_y \end{cases} . \quad (24)$$

It is easy to see that the Casimir operator C (3) does not commute with the Hamiltonian for free particles constrained on the sphere, so it do not represent any invariant for the particles.

IV. CONCLUSION AND DISCUSSION

After analyzing vector operators constructed by directional and angular momentum operator, we find that they can form Lie algebras of type $gl(2, c)$ that can be decomposed into three cases but two algebraic types as $u(1, 1) \otimes u(1)$ and $su(1, 1)$. The unitary irreducible representation for each of these algebras $u(1, 1) \otimes u(1)$ and $su(1, 1)$ can be obtained by standard manner [8–11]. Also from the manner, we can construct the coherent states based on these algebras [5–8], but the relation between these states and the motion of the particle constrained on the sphere is an intricate problem [4–7]. We raise but leave this issue open for further debate and exploration.

-
- [1] Andrews M 1979 *Am. J. Phys.* **47** 274-277
 - [2] Sakurai J J 1994 *Modern Quantum Mechanics* (New York: Addison-Wesley)
 - [3] Ka X L 2001 *Advanced Quantum Mechanics*, 2nd ed. (Beijing: Higher Education) pp.76-77, 102-103
 - [4] Kowalski K and Rembieliński J 2000 *J. Phys. A: Math. Gen.* **33** 6035, and 2002 *J. Math. Phys.* **42** 4238-4147
 - [5] Hall B C and Mitchell J J 2002 *Coherent states on spheres*, *J. Math. Phys.* **43** 1211–1126
 - [6] Fakhri H and Sadeghi J 2004 *International Journal of Theoretical Physics*. **43** 457-476
 - [7] Fakhri H and Chenaghloou A 2007 *J. Phys. A: Math. Theor.* **40** 5511–5523
 - [8] Iachello F 2006 *Lie Algebras and Applications, Lecture Notes in Physics* Vol. 708 (Berlin: Springer)
 - [9] Dong S H 2007 *Factorization Method in Quantum Mechanics, Fundamental Theories of Physics* Vol. 150 (Dordrecht: Springer)
 - [10] Perelemov A 1986 *Generalized Coherent States and Their Applications*. (Berlin: Springer)
 - [11] Klauder J R and Skagerstam B S 1985 *Coherent States—Applications in Physics and Mathematical Physics*. (Singapore: World Scientific)